

Journal of Anatomical Sciences Email: anatomicaljournal@gmail.com

J. Anat Sci 13(2)

Histomorphological and histomorphometric changes in the testes of male Wistar rats following prepubertal exposure to caffeine

<sup>1</sup>Adeyemi DO, <sup>2</sup>Awoniran PO, <sup>1</sup>Orumana OM and <sup>2</sup>Jolayemi KA

<sup>1</sup> Department of Anatomy and Cell Biology, Faculty of Basic Medical Sciences, College of Health Science, Obafemi Awolowo University, Ile-Ife, Nigeria.

<sup>2</sup> Department of Human Anatomy, Faculty of Basic Medical Sciences, Redeemer's University, Ede, Osun State, Nigeria.

**Corresponding author:** Adeyemi DO E-mail: david.adeyemi@oauife.edu.ng; +2348028101028

## ABSTRACT

This experiment was designed to study the effects of caffeine on the histomorphology of the testes of prepubertal male Wistar rats. Prepubertal Wistar rats (50 - 60 g) were randomly divided into three treatment groups (B, C and D) and control (A) of ten rats each. Group A rats received distilled water while Groups B, C, and D rats were respectively given daily dose of 50 mg/kg, 100 mg/kg, and 200 mg/kg of caffeine dissolved in distilled water for 14 days by gavage. Five rats from each group were sacrificed 24 hours after the last caffeine administration while the remaining rats were monitored for another 21 days for them to attain sexual maturity. The rats were sacrificed under diethyl ether anesthesia and the testes were fixed by immersion in Bouin's fluid. The tissues were processed for histomorphological and histomorphometric assessment. Data obtained were analyzed using one way ANOVA followed by Dunnett post hoc test for comparison with control. Alpha level was set at 0.05. Results showed that treatment with caffeine caused a significant decrease in testicular weight and a marked cellular degeneration and depletion of germ cells in the testes of rats in group D. The height of the germinal epithelium was significantly reduced when compared with control. This study concluded that prepubertal exposure of Wistar rats to caffeine reduced testicular growth and deleteriously alters testicular morphology.

**Key words:** Caffeine, Testes, Germinal epithelium, Histomorphology, Histomorphometry, Prepubertal Wistar rats.

### **INTRODUCTION**

Caffeine is a pharmacologically active component of many foods, beverages, dietary supplements, and drugs <sup>[1]</sup>. Being most widely consumed psychoactive drug in the world, its consumption remains largely unregulated and legal in nearly all parts of the world. Unlike most other psychoactive substances. Consumption of high caffeinecontaining energy drinks by the adolescent population has increased markedly in recent years due to the risk-taking behavior of adolescents and their lack of education on the negative effects of caffeine <sup>[2, 3, 4]</sup>. There is growing concern about a decrease in male reproductive health, because a number of laboratory animal studies and human cases have pointed to toxic effects of caffeine on the male reproductive system <sup>[4, 5, 6, 7, 8]</sup>. For instance, several studies on effects of caffeine on testicular morphology and function in men and adult animals have been documented <sup>[5, 9, 10, 11]</sup>. In addition, a number of studies suggest that prenatal and lactational caffeine exposure impairs male gonadal development and thus later gonadal function <sup>[4, 7]</sup>. Since the adolescent gonad is not fully mature either anatomically or histochemically <sup>[12]</sup>, it would not be surprising if chronic exposure to caffeine during this vulnerable period were to interfere with normal gonad maturation. Though detrimental effects of caffeine on the development of the reproductive system during prenatal period have been widely demonstrated, data on the effect on the reproductive system during the prepubertal period are relatively sparse and conflicting. Considering that puberty is a period of rapid development of reproductive capacity, its vulnerability to insults seems to be greater than that in adults <sup>[13]</sup>. This study therefore aimed at investigate the effects of caffeine on morphology, morphometry as well as the parameters of growth and maturation of the testis in prepubertal male rats throughout most of their rapid growth period.

#### MATERIALS AND METHODS

Animals and Treatment: Prepubertal male (4 weeks) Wistar rats (50 - 65 g) were obtained from the Animal House of the College of Health Sciences. Obafemi Awolowo University, Ile–Ife, Nigeria. They were maintained under the condition of natural light and dark cycles, temperature and humidity. Standard laboratory rat chow (Ace feed, Osogbo) and clean water were provided daily to the rats throughout the period of experiment. Ethical clearance was obtained from Health Research and Ethics Committee (HREC) of Institute of Public Health, Obafemi Awolowo University and the guidelines for care and use of animals were strictly followed.

**Experimental Design:** The rats were randomly assigned into four groups of ten rats each (n = 10). The four groups are groups A, B, C and D. Rats in Group A were the normal control and they received distilled water orally for 14 days while Groups B, C, and D received 50 mg/kg, 100 mg/kg and 200 mg/kg graded doses of caffeine (Sigma-Aldrich, USA) respectively were administered orally for a period of 14 days. Twenty-four hours after the last administration, five rats from each group were anaesthetized under diethyl ether and their testes were excised. The remaining rats were monitored for another 21 days to attain sexual maturity after which they were also sacrificed.

**Organ weight:** At the end of the experiment, testes from each rat were excised, trimmed free of fats and weighed on a top loader digital weighing balance (Camry). Relative testicular weight was defined as percentage ratio of testes weight to final body weight. The formula below was used to calculate the organ weight.

*Relative organ weight = organ weight x 100\%* 

final body weight

Histological Procedure: The testes were fixed in fresh Bouin's fluid and tissue processing was done via paraffin wax embedding method of Drury and Wallington <sup>[14]</sup>. The embedded tissues were sectioned on a rotary microtome (Leica RM 2125 RTS) and 5 um thick sections were obtained and stained with Hematoxylin and Eosin for demonstration the of general histoarchitecture. The stained sections were examined under a Leica DM 750 microscope connected to a computerized digital camera system (Leica ICC50) and digital photomicrographs were taken.

#### Histomorphometry:

All

histomorphometric evaluations were performed by the same trained, calibrated, and blinded examiner using an image analysis system (Image J Version 1.4.3.67) to analyze digital photomicrograph taken at a final magnification of 100x. Four serial sections were traced for each testis, and eight measurements per section were made of the number of seminiferous tubules within two defined regions at 100-fold magnification; these measurements were computed to obtain a mean value per animal. The seminiferous tubular epithelia height, seminiferous tubular diameter and luminal seminiferous diameter were measured while the cross-sectional area and volume of the seminiferous tubules were calculated. For analyzing the diameters of the seminiferous tubules, we defined regions that had round or nearly round cross sections of tubules as possible. Then, the longest diameter  $(D_1)$  and the perpendicular diameter (D<sub>2</sub>) of each cross section of tubules were measured; each seminiferous tubular diameter D was calculated as  $\sqrt{D_1D_2}$ .

**Statistical Analysis:** Results were expressed as Mean  $\pm$  S.E.M and data were analyzed using one-way analysis of variance (ANOVA) followed by Dunnett post hoc test for comparison with control. Alpha level was set at 0.05. All statistical analysis was done using GraphPad Prism5 (5.0.3.0, GraphPad Inc.).

# RESULTS

**Organ Weight.** There was a significant decrease in relative testicular weight of treated rats in group D (p < 0.05) when compared with control (Fig. 1). There was no difference in testicular weight between groups A, B, C and D after the recovery period (Fig. 2).



Figure 1: Effects of caffeine on the relative weights of the testes. Values are expressed as the mean  $\pm$  SEM. \* *P*<0.05 vs control.



Figure 2: Effects of caffeine on the relative weights of the testes after 21 days recovery. Values are expressed as the mean  $\pm$  SEM.

**Histopathological Results:** Histological examination of haematoxylin-eosin -stained sections revealed that the rats which received the highest dose (group D), had morphological features of histopathological damages, and these are presented by cellular degeneration resulting in depleted germ cells, a disorganized seminiferous tubular epithelium and presence of pyknotic nuclei (Fig. 3 and 4). However, these features observed in group D rats were reversed after the 3-week recovery period (Fig. 5 and 6). Histological examination of the testes of treated groups B and C rats showed no obvious injury to the seminiferous tubules, spermatogonia cells and the Leydig cells scattered in-between them in the interstitial spaces. They reveal instead seminiferous tubules having normal testicular morphology, primary spermatocytes showing no degenerative changes which showed spermatogenesis was not disturbed (Figs. 5 and 6).



Figure 3: Photomicrographs of sections of testes of prepubertal Wistar rats subjected to H & E stain after treatment. Note in slides A, B, C intact seminiferous tubules showing its lumen (Lu), interstitial cells of Leydig (blue arrows) within the interstitial space (Is) spermatogonia (red arrows), primary spermatocytes (green arrows). Also note D with smaller sizes of seminiferous tubules. Scale bars - 50µm.



Figure 4: Photomicrograph of sections of testes of prepubertal Wistar rats subjected to H & E stain. In slides A, B, and C, note the seminiferous tubule with a regular and thin basement membrane (yellow arrow), lumen (Lu), interstitial spaces (Is) containing interstitial cells (blue arrows), spermatogonia (red arrows), primary spermatocytes (green arrows), Sertoli cells (block arrows). Also note in D germinal epithelium with scanty immature spermatogenic cells. Scale bar  $= 20\mu m$ .



Figure 5: Photomicrographs of sections of testes of prepubertal Wistar rats subjected to H & E stain following recovery. In slides A, B, C, D note the seminiferous tubules showing lumen (Lu), Interstitial cells of Leydig (blue arrows) within the interstitial space (Is), spermatogonia (red arrows) and primary spermatocytes (green arrows). Also note the marked recovery in D. Scale bars - 50µm.



Figure 6: Photomicrograph of sections of testes of prepubertal Wistar rats subjected to H & E stain following recovery. In slides A, B, C, and D note the seminiferous tubule with regular and thin basement membrane (yellow arrow), lumen (Lu) containing mature spermatozoa, interstitial spaces (Is) consisting of interstitial cells of Leydig (blue arrows), spermatogonia (red arrows), spermatocytes (green arrows), Sertoli cells (block arrows), round spermatids (black arrows), flagellum of elongated spermatids within the lumen (arrow heads). Scale bar = 20μm

**Histomorphometry:** Histomorphometric analysis of the seminiferous tubular diameter presented nearly no differences (p > 0.05) between control group (A) and each of the other treated groups (B and C), while group D showed a significant reduction (p < 0.05) in seminiferous tubular diameter in comparison to group A and a significant reduction (p < 0.05) in seminiferous epithelial height in comparison to groups A and B (Table 1). The seminiferous luminal diameter was significantly increased in group D when compared with the other groups (Table 1). At recovery, significant decrease was observed in seminiferous epithelia height of groups C and D when compared with control. A significant decrease in the seminiferous tubular diameter of groups B, C and D was also observed when compared with control while seminiferous luminal diameter of group D increased significantly (p < 0.05) when compared with control (Table 2). In addition, the cross-sectional area of the seminiferous tubule and the volume of the seminiferous tubules significantly decreased (p < 0.05) in groups B, C, and D compared to the control group A.

Groups	Seminiferous	Seminiferous tubular	Seminiferous I	Luminal Cross-sectional area of	Volume of Seminiferous
	tubular diameter	epithelia height	diameter (µm)	seminiferous tubule	tubules (μm <sup>3</sup> )
	(µm)	(µm)		$(mm^2)$	
Gp 1A <sub>1</sub>	$249.8\pm10.92$	$0.229\pm0.019$	$76.02\pm3.377$	$49.02 \text{ x } 10^3 \pm 9.4 \text{ x } 10^1$	$8.16 \ge 10^6 \pm 6.8 \ge 10^3$
Gp 1B <sub>1</sub>	$222.8\pm19.62$	$0.179\pm0.013$	$90.6\pm9.027$	$38.99 \ge 10^3 \pm 30.2 \ge 10^1$	$5.79 \ge 10^6 \pm 3.9 \ge 10^4$
Gp 1C <sub>1</sub>	221.1 ±11.13	$0.180\pm0.003$	$86.6\pm5.086$	$34.58 \ge 10^3 \pm 30.5 \ge 10^1$	$4.84 \ x \ 10^6 \pm 4.0 \ x \ 10^4$
Gp 1D <sub>1</sub>	$209.8\pm19.72\texttt{*}$	$0.155 \pm 0.025^{*}$	$102.3 \pm 8.421*$	$38.40 \ge 10^3 \pm 9.7 \ge 10^1$	$5.66 \ge 10^6 \pm 7.2 \ge 10^3$

Table 1:Effects of caffeine on the morphometry of the testis following treatment with caffeine.

Values are presented as Mean  $\pm$  SEM (n=5).\* = Significantly different from Control (p<0.05)

**Table 2**:
 Effects of caffeine on the morphometry of the testis following recovery from treatment with caffeine.

Groups	Seminiferous	Seminiferous	Seminiferous	Cross-sectional area of	Volume of Seminiferous
	tubular diameter	tubular epithelia	Luminal diameter	seminiferous tubule	tubules (μm <sup>3</sup> )
	(µm)	height (µm)	(μm)	(mm <sup>2</sup> )	
Gp 1A <sub>2</sub>	$320.8\pm22.43$	$0.281\pm0.009$	$122.2 \pm 5.975$	$80.84 \text{ x } 10^3 \pm 4.0 \text{ x } 10^2$	$17.29 \text{ x } 10^6 \pm 5.9 \text{ x } 10^4$
Gp 1B <sub>2</sub>	$274.0 \pm 10.89*$	$0.239\pm0.005$	$115.8\pm4.102$	58.97 x $10^3 \pm 9.3$ x $10^{1*}$	$10.77 \ge 10^6 \pm 6.8 \ge 10^{3*}$
$Gp 1C_2$	$250.1 \pm 6.912*$	$0.194\pm0.007\texttt{*}$	$121.7\pm4.344$	$49.13 \ x \ 10^3 \pm 3.8 \ x \ 10^{1*}$	$8.19 \ge 10^6 \pm 1.7 \ge 10^{3*}$
Gp 1D <sub>2</sub>	$252.5 \pm 13.41*$	$0.157 \pm 0.028*$	$144.6 \pm 12.820 *$	$50.08 \ x \ 10^3 \pm 1.4 \ x \ 10^{2*}$	$8.43 \ x \ 10^6 \pm 1.3 \ x \ 10^{4*}$

Values are presented as Mean  $\pm$  SEM (n=5). \* = Significantly different from Control (p<0.05)

# DISCUSSION

Morphological assessment of the testes of rats exposed to caffeine in this study revealed that caffeine intake caused a significant reduction on in testicular weight. degeneration and marked cellular a depletion of germ cells, a significant height of the germinal decrease in epithelium, seminiferous luminal diameter, seminiferous tubular diameter. cross sectional area and volume of seminiferous tubules of Wistar rats. This observation is in line with result of previous studies in which caffeine toxicity was reported to be associated with a decrease in germ cell proliferation, hence a reduced number of differentiating germ cells <sup>[11]</sup>, as the weight of the testis is directly proportional to the mass of the differentiated spermatogenic cells <sup>[15]</sup>. However, this morphological disruption appears to be reversible after the period of recovery. Oluwole et al. [16] opined that the decrease in organ weight caused by caffeine, could be due to the ability of caffeine to cause lipolysis. This resulted from inhibition of cyclic nucleotide phosphodiesterase that leads to accumulation of cvclic AMP in tissues which then activates hormone-sensitive lipase to promotes lypolysis <sup>[17]</sup>. The decrease in organ weight, may also have resulted from diuresis which hinders reabsorption of salts and water, or increased metabolism <sup>[18]</sup>. This study is also in line with the observations of Dorostghoal et al. <sup>[4]</sup> who reported that prenatal caffeine exposure brought about a dose-related reduction in the testis weight of male offspring.

The examination of the histological sections of group D showed that caffeine induced deleterious effects on testicular architecture which could have resulted from interference with the testicular micro-environment thereby resulting in the disruption of the initiation and maintenance of the spermatogenesis. A reduction was observed in seminiferous tubular epithelia height of the group administered with the highest dose reveals shows caffeine has negative effect on testicular morphology. Pollard et al. <sup>[6]</sup> had earlier reported that exposure of male Wistar rats to 30 mg/kg per day of caffeine, given by gavage for 15 to 38 consecutive days led to breakdown of the germinal epithelium. The seminiferous tubular diameter was observed to be decreased in all the animals after the recoverv period. The decrease in seminiferous tubular diameter may be due to the fact that caffeine interferes with cell division which leads to a reduced number of spermatogenic cells. Dorostghoal et al. [4], reported a decrease in the diameter of seminiferous tubules of the offspring of rats given caffeine during the prepubertal. postpubertal, and adulthood periods. Pollard et al. [6] also showed that caffeine when given prenatally, inhibited differentiation of the seminiferous cords. Daily administration of 30 or 60 mg/kg caffeine to mature male rabbits for four consecutive weeks caused a decrease in the sizes of the seminiferous tubules and inhibited spermatogenesis <sup>[19]</sup>.

# CONCLUSION

This study concluded that prepubertal exposure of Wistar rats to caffeine reduced testicular growth and predisposed the seminiferous tubules of growing male Wistar rats to deleterious structural changes which were reversible over a recovery period.

### **CONFLICT OF INTEREST**

The research received no funding from any profit or non-profit making organization. Authors therefore, declare that there are no potential conflicts of interest.

### REFERENCES

- 1 Wikoff D, Welsh BT, Henderson R, Brorby GP, Britt J, Myers E, Goldberger J. Lieberman HR. O'Brien C. Peck J. Tenenbein M. Weaver C, Harvey S, Urban J, and Doepker C. Systematic review of the potential adverse effects of caffeine consumption healthy in adults. pregnant women. adolescents. and children, Food and Chemical Toxicology, 2017; 109 (1): 585 - 648 https://doi.org/10.1016/j.fct.2017.04.0 02.
- Reissig C.J., Strain E. C., and Griffiths R. R. Caffeinated energy drinks—a growing problem, Drug and Alcohol Dependence. 2009; 99 (1–3): 1–10.
- 3. Seifert S. M., Schaechter J. L., Hershorin E. R., and Lipshultz S. E. Health effects of energy drinks on children, adolescents, and young adults. Pediatrics; 2011; 127 (3): 511–528.
- 4. Dorostghoal M., Erfani-Majd N., and Nooraei P. Maternal caffeine consumption has irreversible effects on reproductive parameters and fertility in male offspring rats. *Clin. Exp. Reprod. Med.* 2012; 39: 144– 152. doi: 10.5653/cerm.2012.39.4.144
- 5. Weinberger MA, Friedman L, Farber TM, Moreland FM, Peters EL, Gilmore CE. and Khan MA. Testicular atrophy and impaired spermatogenesis in rats fed high levels of the methylxanthines caffeine. theobromine. or theophylline. Journal of Environmental Pathology and Toxicology, 1978; 1(5):669-688.
- 6. Pollard I, Williamson S., and Magre S. Influence of caffeine administered during pregnancy on the early differentiation of fetal rat ovaries and testes, Journal of Developmental Physiology, 1990; 13 (2): 59–65.

- Ramlau-Hansen C. H., Thulstrup A. M., Bonde J. P., Olsen J, and Bech B. H. Semen quality according to prenatal coffee and present caffeine exposure: two decades of follow-up of a pregnancy cohort, Human Reproduction, 2008; 23 (12): 2799– 2805. doi: 10.1093/humrep/den331
- 8. Tinwell H., Colombel S., Blanck O., and Bars R. The screening of everyday life chemicals in validated assays targeting the pituitary-gonadal axis, Regulatory Toxicology and Pharmacology, 2013; 66 (2): 184– 196.
- Nawrot P., Jordan S., Eastwood J., Rotstein J., Hugenholtz A., Feeley M. Effects of caffeine on human health. *Food Addit. Contam.* 2003; 20: 1–30.
- 10. Park, M.; Choi, Y.; Choi, H.; Yim, J.Y. and Roh, J. High Doses of Caffeine during the Peripubertal Period in the Rat Impair the Growth and Function of the Testis. *Int. J. Endocrinol.* 2015; Pp368-475.
- 11. Bae, J.; Choi, H.; Choi, Y. Roh, J. Dose- and time-related effects of caffeine on the testis in immature male rats. *Exp. Anim.*, 2015; 66:29-39.
- August G. P., Grumbach M. M., Kaplan S. L. Hormonal changes in puberty: III. Correlation of plasma testosterone, LH, FSH, Testicular size, and bone age with male pubertal development. The Journal of Clinical Endocrinology and Metabolism, 1972; 34 (2): 319–326.
- 13. Sharpe R. Endocrinology and paracrinology of the testis. *Physiology and toxicology of male reproduction*. 1988; Pp 71–102.
- Drury, R.A., Wallington, E.A. Carleton's Histological Techniques.
   5th Edition. New York: Oxford University Press. 1980; Pp 44 – 134.
- 15. Boockfor, F.R. and Blake, C.A. Chronic administration of 4-tertoctylphenol to adult male rats causes

shrinkage of the testes and male accessory sex organs, disrupts spermatogenesis, and increases the incidence of sperm deformities. *Biol. Reprod.*, 1997; 57:267–277.

- Oluwole, O.F., 16. Salami, S.A., Ogunwole. E. and Raii. Y. Laboratory for Reproductive Physiology Developmental and Programming, Implication of caffeine consumption and recovery on the reproductive functions of adult male Wistar rats. J Basic Clin Physiol Pharmacol., 2016; 1-8.
- Hursel, R, and Westerterp-Plantenga, M.S. Thermogenic ingredients and body weight regulation. *Int J Obes.*, 2010; 34:659–69.

- Huang, Z.L., Qu, W.M., Eguchi, N., Chen, J.N., Schwarzschild, M.A. and Fredholm, B.B. Adenosine A2A, but not A1, receptors mediate the arousal effect of caffeine. *Nat Neurosci.*, 2005; 8:858–9.
- 19. Ezzat, A.R. and el-Gohary, Z.M. Hormonal and histological effects of chronic caffeine administration on the pituitary-gonadal and pituitaryadrenocortical axes in male rabbits. *Functional and developmental morphology.*, 1994; 4:45-50.